

**Recommended dosing regimens of Sabvixin®
in patients with normal renal function**

Indication	Daily dose regimen	Way of administration	Duration of treatment
Bacterial meningitis	Adults and children under 2 years old: 5 mg once a day during 3-4 days, then 5 mg once every other day for at least 2 weeks after cultures of the cerebrospinal fluid are negative and sugar content has returned to normal Children under 2 years old: 2 mg once a day during 3-4 days, then 2 mg once every other day for at least 2 weeks after cultures of the cerebrospinal fluid are negative and sugar content has returned to normal	intrathecally	14-18 days
Nosocomial pneumonia, including ventilator-associated pneumonia	Adults: 1.2 mg/kg every 12 hours Children ¹ : 1.2 mg/kg every 12 hours	IV	10-14 days
Skin and soft tissues infections caused by gram-negative bacteria	Adults: 1.2 mg/kg every 12 hours <i>or</i> 1 mg/kg every 8 hours Children ¹ : 1.2 mg/kg every 12 hours	IV IM	10-14 days
Burn wound infection caused by <i>P. aeruginosa</i> or <i>Acinetobacter</i> spp.	Adults: 1.2 mg/kg every 12 hours <i>or</i> 1 mg/kg every 8 hours	IV IM	10-14 days

Gram-negative sepsis	Adults: 1.2 mg/kg every 12 hours Children ¹ : 1.2 mg/kg every 12 hours	IV	10-14 days
Eye infections caused by <i>P. aeruginosa</i> (keratitis, conjunctivitis, corneal ulcer)	1-3 drops of 0.1-0.25% solution hourly ² 1-5 mg once a day	Eye drops Subconjunctival injection	5-7 days
Selective decontamination of the digestive tract in mechanically ventilated patients	10 ml of aqueous suspension containing 100 mg of polymyxin B, 80 mg of gentamicin or tobramycin, and 500 mg of amphotericin B every 6 hours	Through a nasogastric tube	During the period of mechanical ventilation and 48 hours after extubation
Selective decontamination of the digestive tract in patients with severe acute pancreatitis	10 ml of aqueous suspension containing 100 mg of polymyxin B, 80 mg of gentamicin or tobramycin, and 500 mg of amphotericin B every 6 hours	Orally or through a nasogastric tube	Up to 7 days after the onset of disease

¹ for children over 1-year-old; for children under 1-year-old the daily dose may be increased to 4 mg/kg and divided into 2 intravenous injections with a 12-hour interval, or into 3-4 intramuscular injections with a 6-8-hour interval

² to prepare eye drops dissolve 50mg polymyxin B in 20 to 50 mL sterile water for injection or sodium chloride injection for a 0.1%-0.25% concentration; solutions of polymyxin B can be stored for 72 hours at the temperature $5 \pm 3^{\circ}\text{C}$.