

**Recommended dosing regimens of Ampicillin + sulbactam,
produced by Kraspharma OJSC, in patients with normal renal function**

Indication	Daily dose regimen	Way of administration	Duration of treatment
Secondary peritonitis (including appendicular peritonitis) Diverticulitis Perirectal abscess	Adults: 3 g every 6 hours, or 4.5 g every 8 hours (maximum daily dose of sulbactam is 4 g) Children ¹ : 200-300 mg/kg/day, divide into 4 injections	IV IV	7-10 days
Pleural empyema Lung abscess	Adults: 3 g every 6 hours, or 4.5 g every 8 hours (maximum daily dose of sulbactam is 4 g) Children ¹ : 200-300 mg/kg/day, divide into 4 injections +/- gentamicin 5 mg/kg/day, divide into 2 injections	IV or IM IV or IM IV or IM	21-42 days, step- down therapy is possible
Amnionitis Endometritis Septic abortion	Adults: 3 g every 6 hours (maximum daily dose of sulbactam is 4 g)	IV or IM	7-10 days
Biliary tract infection, including post- operative cholangitis	Adults: 3 g every 6 hours Children ¹ : 150-200 mg/kg/day, divide into 4 injections	IV or IM IV or IM	10-14 days
Orchiepididymitis (men > 35 yrs)	1.5 g every 6 hours	IV or IM	10-14 days

Post-operative wound infections	Adults: 1.5-3 g every 6 hours	IV or IM	7-10 days
Diabetic foot infections	Children ¹ : 150-300 mg/kg/day, divide into 4 injections	IV or IM	
Bite wounds			
Sepsis	Adults: 3 g every 6 hours (maximum daily dose of sulbactam is 4 g) Children ¹ : 300 mg/kg/day, divide into 4 injections	IV IV	10-14 days
Perioperative antibiotic prophylaxis (in traumatology, obstetrics, gynecology, abdominal & colorectal surgery)	Adults: 1.5-3 g 30 min. prior to incision Children ¹ : 50-75 mg/kg 30 min. prior to incision	IV IV	

¹ – doses for children over 1 year of age