

## Recommended dosing regimens of Cefoperazone, produced by Kraspharma OJSC

Indication	Daily dose regimen	Way of administration	Duration of treatment
Nosocomial pneumonia	<p>Adults: 1-2 g every 8-12 hours +/- amikacin 15-20 mg/kg/day in 1 to 2 divided doses</p> <p>Children: 80 mg/kg/day divided into two doses +/- amikacin 15-20 mg/kg/day in 1 to 2 divided doses</p>	<p>IV or IM</p> <p>IV or IM</p> <p>IV or IM</p> <p>IV or IM</p>	10-14 days
Intra-abdominal infections, including secondary peritonitis	<p>Adults: 2 g every 8-12 hours + metronidazole 0.5 g every 8 hours</p> <p>Children: 80 mg/kg/day divided into two doses + metronidazole 22.5 mg/kg/day in 3 equally divided doses</p> <p>Newborns: 30 mg/kg every 12 hours + metronidazole 7.5-15 mg/kg every 12 hours</p>	<p>IV or IM</p> <p>IV</p> <p>IV or IM</p> <p>IV</p> <p>IV</p> <p>IV</p>	7-10 days
Biliary tract infection (cholangitis, cholecystitis)	<p>Adults: 1-2 g every 8-12 hours + metronidazole 0.5 g every 8 hours</p> <p>Children: 80 mg/kg/day divided into two doses + metronidazole 22.5 mg/kg/day in 3 equally divided doses</p>	<p>IV or IM</p> <p>IV</p> <p>IV or IM</p> <p>IV</p>	7-10 days

Acute pyelonephritis Post-operative urinary tract infection	Adults: 1-2 g every 8-12 hours Children: 80 mg/kg/day divided into two doses	IV or IM  IV or IM	14 days
Pelvic inflammatory disease (PID)	2 g every 12 hours + metronidazole 500 mg every 8 hours +/- doxycycline 100 mg every 12 hours	IV or IM IV or per os  IV or per os	10-14 days
Sepsis (source – abdominal cavity or female reproductive tract), as well as biliary sepsis	Adults: 2-4 g every 8-12 hours + metronidazole 0.5 g every 8 hours Children: 100-200 mg/kg/day divided into two or three doses + metronidazole 22.5 mg/kg/day in 3 equally divided doses Newborns: 35-40 mg/kg every 12 hours + metronidazole 7.5- 15 mg/kg every 12 hours	IV IV  IV  IV IV	10-14 days